

Final Technical Report

USGS Award 02HQGR0039

HIGH RESOLUTION STUDIES OF CALAVERAS FAULT SEISMICITY

P.I. Gregory C. Beroza

Program Element II. Earthquake Physics and Effects

Research supported by the U.S. Geological Survey (USGS), Department of the Interior, under USGS award number 02-HQ-GR-0039. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

Award # 02-HQ-GR-0039

HIGH RESOLUTION STUDIES OF CALAVERAS FAULT SEISMICITY

Gregory C. Beroza
Department of Geophysics
397 Panama Mall
Stanford, CA, 94305-2215

Phone: (650) 723-4958
Fax: (650) 725-7344
beroza@geo.stanford.edu

Technical Abstract

By measuring relative earthquake arrival times using waveform cross correlation and locating earthquakes using the double-difference technique, we are able to reduce hypocentral errors by 1 to 2 orders of magnitude over routine locations for nearly 8000 earthquakes along a 35-km section of the Calaveras Fault. This represents ~92% of all seismicity on this fault from 1984-1998 and includes the rupture zone of the M 6.2 1984 Morgan Hill, California, earthquake. The relocated seismicity forms highly organized structures that were previously obscured by location errors. There are abundant repeating earthquake sequences as well as linear clusters of earthquakes. Large voids in seismicity appear with dimensions of kilometers that have been aseismic over the 30-year time interval, suggesting that these portions of the fault are either locked or creeping. The area of greatest slip in the Morgan Hill main shock coincides with the most prominent of these voids, suggesting that this part of the fault may be locked between large earthquakes. We find that the Calaveras Fault at depth is extremely thin, with an average upper bound on fault zone width of 75 m. Given the location error, however, this width is not resolvably different from zero. The relocations reveal active secondary faults, which we use to solve for the stress field in the immediate vicinity of the Calaveras Fault. We find that the maximum compressive stress is at a high angle, only 13° from fault normal, even to within a few hundred meters of the Calaveras fault, supporting previous interpretations that this fault is weak.

Award # 02-HQ-GR-0039
High Resolution Studies of Calaveras Fault Seismicity

Gregory C. Beroza
Department of Geophysics
397 Panama Mall
Stanford, CA, 94305-2215

Phone: (650) 723-4958
Fax: (650) 725-7344
beroza@geo.stanford.edu

Non-Technical Abstract

By careful measurement of the arrival times of earthquake waves in combination with improved earthquake location techniques, we are able to reduce earthquake location errors from over 1 km to 100 meters or less for nearly 8000 earthquakes along a 35-km section of the Calaveras Fault in California. The improved earthquake locations allow us to discern highly organized structures in the seismicity that were previously obscured. We find many sequences of repeating earthquake sequences, linear clusters of earthquakes, large voids in seismicity on the Calaveras fault. The relocations also reveal active secondary faults that support previous interpretations that the Calaveras fault is weak.

Introduction

The Calaveras Fault (figure 1) is one of the most active branches of the San Andreas system in northern California. In the 20th century there have been a series of moderate magnitude earthquakes some of which have repeated [Oppenheimer et al., 1990]. The largest of these earthquakes was the M 6.2 1984 Morgan Hill earthquake, which appears to have been a repeat of an earthquake in 1911. The long-term slip rate on the Calaveras Fault to the south of where the Hayward Fault splays from it is thought to be 15 ± 3 mm/yr [Working Group on California Earthquake Probabilities (WGCEP), 1999].

In addition to rupture in moderate earthquakes, this section of the Calaveras Fault has been extraordinarily active in microearthquake activity. It is also known to creep with an average rate of ~ 13.5 mm/yr since 1997 [WGCEP, 1999]. Because of the combination of a fast slip rate, the presence of creep, and equivocal paleoseismic evidence for large earthquakes in the past, the potential for future large earthquakes on the central section of the Calaveras Fault remains uncertain [WGCEP, 1999].

Oppenheimer et al. [1990] studied microearthquake activity of the Calaveras Fault in great detail. They found a correspondence between the areas that slipped in moderate earthquakes with areas that were relatively devoid of microearthquake activity. They also found that small earthquakes had a very similar spatial distribution both before and after moderate earthquakes. They proposed that the areas devoid of seismicity were stuck between moderate earthquakes and used this assumption to identify two likely source zones for future moderate earthquakes on the Calaveras Fault.

The geometry of fault zones within the Earth is revealed primarily by seismicity. In this study, we greatly improve the earthquake locations on the Calaveras Fault. Our ability to resolve fault zone structure and hence to address many essential aspects of earthquake behavior, is limited by our ability to obtain precise earthquake locations. While quantitative earthquake location techniques date back to the early years of the 20th century [e.g., Geiger, 1910], our incomplete knowledge of Earth structure and the difficulty of measuring arrival times accurately typically limits the accuracy of earthquake locations to tens of kilometers at teleseismic distances and several kilometers at regional distances even in well instrumented regions like California and Japan.

Uncertainty in earthquake locations can be dramatically reduced by aiming for precision, rather than accuracy, in the form of relative earthquake locations [e.g., Poupinet et al., 1984; Fre'chet, 1985; Deichmann and Garcia-Fernandez, 1992; Got et al., 1994]. In this study, we improve the precision of earthquake locations in two principal ways: (1) removing substantial velocity model error with the double difference approach [Waldhauser and Ellsworth, 2000] and (2) reducing relative arrival time measurement errors by waveform cross correlation [Schaff, 2001]. The combination allows us to reduce relative location errors by 1 to 2 orders of magnitude for the entire data set.

What emerges from the improved earthquake locations is a highly refined picture of seismicity on the Calaveras Fault (figures 2,3,4). Numerous features that were seen as

clouds of seismicity are brought into sharp focus (figure 2). These features include streaks of earthquakes that are elongated in the direction of slip, repeating microearthquakes that recur up to dozens of times, and large areas of little seismicity. We obtain a clearer view of a kilometer-scale compressional fault offset and its geometry at depth. We are also able to resolve secondary fault structures, which were previously suggested by focal mechanisms [Oppenheimer et al., 1988]. Slip on these structures indicate that the maximum compressive stress is at a high angle to the Calaveras Fault implying that it is weak if the stress remains unfavorably oriented on the fault plane. This seems likely, as secondary structures are active to within 100 m of the main fault trace. If the stress rotates into a more favorable orientation, it must do so within several hundred meters of the fault. The improved locations also allow us to constrain the degree of strain localization on the Calaveras Fault. We find a very narrow fault zone width of no more than 75 m at seismogenic depths.




Figure 1. Study area is a 35 km segment of the Calaveras fault near San Jose, California. There were nearly 8000 earthquakes on the fault during the period of study. We used approximately 8,000,000 cross correlation arrival time measurements of *P* and *S* waves to relocate these earthquakes.




Figure 2. Improvement in earthquake locations is demonstrated by comparing the standard catalog locations (left panel) with the improved locations (right panel). The previously chaotic seismicity becomes highly organized after relocation.




Figure 3. Map view of earthquakes before (top) and after (bottom) relocation. The map is oriented such that the trend of the nearly vertical Calaveras fault is horizontal on the figure. The fault discontinuity at 14-18 km along strike is much more clearly defined in the relocations. Many of the off-fault earthquakes above the fault in the figure show up much more clearly in the relocations. These are right-lateral (not conjugate) strike-slip faults and the trend of the microearthquakes is consistent with their focal mechanisms. The more diffuse seismicity shown below the mainshock fault plane defines a clear dipping planar fault in cross section. Here too the reverse faulting focal mechanisms of these events is consistent with the planes of microearthquakes in the relocations. Together with the mainshock fault plane, these combinations of active faults constrain the maximum compressive stress to be nearly normal to the Calaveras Fault. The mainshock fault plane itself is much more sharply defined in the relocations and the width of the fault at depth is not resolvably different from zero.




Figure 4. Side view of earthquakes before (top) and after (bottom) relocation. The earthquakes are plotted as circles with the size of the circle corresponding to that expected for a circular constant stress drop of 3 MPa. Much of the fault is not covered with seismicity and there are many areas of frequently repeating earthquakes. The large circle is the 1984 Morgan Hill California earthquake. This earthquake is large enough that there are independent determinations of the slip distribution from strong motion data [Beroza and Spudich, 1988]. These indicate that most of the slip occurred in areas devoid of microearthquake activity, which in turn suggests that these areas are locked between large earthquakes. The streaks of seismicity are roughly aligned with the slip direction. A temporary deepening of the deepest earthquakes from -1 to -4 km along strike (not shown) following the 1984 Morgan Hill earthquake is consistent with the notion that a strain-rate dependent rheology controls the depth of the seismic to aseismic transition.

A complete description of our results for the Calaveras Fault was published in the Journal of Geophysical Research as:

Schaff, D. P., G. H. R. Bokelmann, G. C. Beroza, F. Waldhauser, and W. L. Ellsworth, High resolution image of Calaveras Fault seismicity, *J. Geophys. Res.*, **107** (B9), 2186, doi:10.1029/2001JB000633, 2002.

A description of the cross correlation measurement techniques that were used to develop these locations was published in the Bulletin of the Seismological Society of America as:

Schaff, D.P., G.H.R. Bokelmann, W. L. Ellsworth, E. Zanker, F. Waldhauser, and G. C. Beroza, Optimizing correlation techniques for improved earthquake location, *Bull. Seismol. Soc. Am.*, **94**, 705-721, 2004.

References

- Angelier, J., Determination of the mean principal directions of stresses for a given fault population, *Tectonophysics*, 56, T17–T26, 1979.
- Bakun, W. H., M. M. Clark, R. S. Cockerham, W. L. Ellsworth, A. G. Lindh, W. H. Prescott, A. F. Shakal, and P. Spudich, The 1984 Morgan Hill, California, earthquake, *Science*, 225, 288–291, 1984.
- Bakun, W. H., G. C. P. King, and R. S. Cockerham, Seismic slip, aseismic slip, and the mechanics of repeating earthquakes on the Calaveras Fault, California, in *Earthquake Source Mechanics*, *Geophys. Monogr. Ser.*, vol. 37, edited by S. Das, J. Boatright, and C. H. Scholz, pp. 195–207, AGU, Washington, D. C., 1986.
- Beeler, N. M., D. A. Lockner, and S. H. Hickman, A simple stick-slip and creep slip model for repeating earthquakes and its implication for microearthquakes at Parkfield, *Bull. Seismol. Soc. Am.*, 91, 1797–1804, 2001.
- Beroza, G. C., and P. Spudich, Linearized inversion for fault rupture behavior; application to the 1984 Morgan Hill, California, earthquake, *J. Geophys. Res.*, 93, 6275–6296, 1988.
- Byerlee, J. D., Friction of rocks, *Pure Appl. Geophys.*, 116, 615–629, 1978.
- Deichmann, N., and M. Garcia-Fernandez, Rupture geometry from high precision relative hypocentre locations of microearthquake ruptures, *Geophys. J. Int.*, 110, 501–517, 1992.
- Ellsworth, W. L., Characteristic earthquakes and long-term earthquake forecasts: Implications of central California seismicity, in *Urban Disaster Mitigation*, edited by F. Y. Cheng and M. S. Sheu, pp. 1–14, Elsevier Sci., New York, 1995.
- Felzer, K. R., and G. C. Beroza, Deep structure of a fault discontinuity, *Geophys. Res. Lett.*, 26, 2121–2124, 1999.
- Fre'chet, J., *Sismogene'se et doublets sismiques, the'se d'etat*, 206 pp., Univ. Sci. et Me'd. de Grenoble, Grenoble, France, 1985.

- Geiger, L., Herdbestimmung bei Erdbeben aus den Ankunftszeiten, *K. Ges. Wiss. Goett.*, 4, 331–349, 1910.
- Gillard, D., A. M. Rubin, and P. Okubo, Highly concentrated seismicity caused by deformation of Kilauea's deep magma system, *Nature*, 384, 343–346, 1996.
- Got, J.-L., J. Fre'chet, and F. W. Klein, Deep fault plane geometry inferred from multiplet relative relocation beneath the south flank of Kilauea, *J. Geophys. Res.*, 99, 15,375–15,386, 1994.
- Hardebeck, J. L., and E. Hauksson, Role of fluids in faulting inferred from stress field signatures, *Science*, 285, 236–239, 1999.
- Hartzell, S. H., and T. H. Heaton, Rupture history of the 1984 Morgan Hill, California, earthquake from the inversion of strong motion records, *Bull. Seismol. Soc. Am.*, 76, 649–674, 1986.
- Li, Y.-G., P. C. Leary, K. Aki, and P. Malin, Seismic trapped modes in the Oroville and San Andreas Fault zones, *Science*, 249, 763–766, 1990.
- Li, Y.-G., J. E. Vidale, K. Aki, C. J. Marone, and W. H. K. Lee, Fine structure of the Landers fault zone; segmentation and the rupture process, *Science*, 265, 367–370, 1994.
- Li, Y.-G., K. Aki, and F. L. Vernon, San Jacinto fault zone guided waves; a discrimination for recently active fault strands near Anza, California, *J. Geophys. Res.*, 102, 11,689–11,701, 1997.
- Li, Y.-G., K. Aki, J. E. Vidale, and M. G. Alvarez, A delineation of the Nojima Fault ruptured in the M 7.2 Kobe, Japan, earthquake of 1995 using fault zone trapped waves, *J. Geophys. Res.*, 103, 74,247–74,263, 1998.
- Mendoza, C., and S. H. Hartzell, Aftershock patterns and main shock faulting, *Bull. Seismol. Soc. Am.*, 78, 1438–1449, 1988.
- Michael, A., Effects of three-dimensional velocity structure on the seismicity of the 1984 Morgan Hill, California, aftershock sequence, *Bull. Seismol. Soc. Am.*, 78, 1199–1221, 1988.
- Nadeau, R. M., and L. R. Johnson, Seismological studies at Parkfield, VI, Moment release rates and estimates of source parameters for small repeating earthquakes, *Bull. Seismol. Soc. Am.*, 88, 790–814, 1998.
- Nadeau, R. M., and T. V. McEvilly, Fault slip rates at depth from recurrence intervals of repeating microearthquakes, *Science*, 285, 718–721, 1999.
- Oppenheimer, D. H., P. A. Reasenber, and R. W. Simpson, Fault plane solutions for the 1984 Morgan Hill, California, earthquake sequence; evidence for the state of stress on the Calaveras Fault, *J. Geophys. Res.*, 93, 9007–9026, 1988.
- Oppenheimer, D. H., W. H. Bakun, and A. G. Lindh, Slip partitioning of the Calaveras Fault, California, and prospects for future earthquakes, *J. Geophys. Res.*, 95, 8483–8498, 1990.
- Poupinet, G., W. L. Ellsworth, and J. Fre'chet, Monitoring velocity variations in the crust using earthquake doublets: An application to the Calaveras Fault, California, *J. Geophys. Res.*, 89, 5719–5731, 1984.
- Prescott, W. H., N. E. King, and G. Guohua, Preseismic, coseismic, and postseismic deformation associated with the 1984 Morgan Hill, California, earthquake, in *The 1984 Morgan Hill, CA earthquake*, edited by J. H. Bennett and R. W. Sherburne, Spec. Publ. Calif. Div. Mines Geol., 68, 137–148, 1984.

- Radbruch-Hall, D. H., Map showing recently active breaks along the Hayward Fault zone and the southern part of the Calaveras Fault zone, California, U. S. Geol. Surv. Misc. Invest., I-813, 1974.
- Rubin, A. M., Aftershocks of microearthquakes as probes of the mechanics of rupture, *J. Geophys. Res.*, 107(B7), 2142, doi:10.1029/2001JB000496, 2002.
- Rubin, A. M., D. Gillard, and J.-L. Got, Streaks of microearthquakes along creeping faults, *Nature*, 400, 635– 641, 1999.
- Sammis, C. G., and J. R. Rice, Repeating earthquakes as low-stress drop events at a border between locked and creeping fault patches, *Bull. Seismol. Soc. Am.*, 91, 532–537, 2001.
- Schaff, D. P., 4D high resolution seismology: Repeating events and large scale relocation, Ph.D. thesis, 115 pp., Stanford Univ., Stanford, Calif., 2001.
- Schaff, D. P., G. C. Beroza, and B. E. Shaw, Postseismic response of repeating aftershocks, *Geophys. Res. Lett.*, 25, 4549– 4552, 1998.
- Sibson, R. H., Roughness at the base of the seismogenic zone: Contributing factors, *J. Geophys. Res.*, 89, 5791– 5799, 1984.
- Townend, J., and M. D. Zoback, Implications of earthquake focal mechanisms for the frictional strength of the San Andreas Fault system. in *The Nature and Tectonic Significance of Fault Zone Weakening*, edited by R. E. Holdsworth et al., *Geol. Soc. Spec. Publ.*, 186, 13– 21, 2001.
- Waldhauser, F., A computer program to compute double-difference hypocenter locations, U.S. Geol. Surv. Open File Rep., 01-113, 25 pp., 2001.
- Waldhauser, F., and W. L. Ellsworth, A double-difference earthquake location algorithm: Method and application to the northern Hayward Fault, California, *Bull. Seismol. Soc. Am.*, 90, 1353– 1368, 2000.
- Waldhauser, F., and W. L. Ellsworth, Fault structure and mechanics of the Hayward Fault, California, from double-difference earthquake locations, *J. Geophys. Res.*, 107(B3), 2054, doi:10.1029/2000JB000084, 2002.
- Waldhauser, F., W. L. Ellsworth, and A. Cole, Slip-parallel seismic lineations along the northern Hayward Fault, California, *Geophys. Res. Lett.*, 26, 3525– 3528, 1999.
- Working Group on California Earthquake Probabilities (WGCEP), Earthquake probabilities in the San Francisco Bay region: 2000 to 2030—A summary of findings, online version 1.0, U.S. Geol. Surv. Open file Rep., 99-517, 60 pp., 1999.
- Zoback, M. D., et al., New evidence on the state of stress of the San Andreas Fault system, *Science*, 238, 1105– 1111, 1987.